

# Genetic analysis of starch paste viscosity parameters in glutinous rice (*Oryza sativa* L.)

Chang-Jie Yan · Zhi-Xi Tian · Yu-Wei Fang · Ya-Chun Yang · Jian Li ·  
Sheng-Yuan Zeng · Shi-Liang Gu · Chen-Wu Xu · Shu-Zhu Tang ·  
Ming-Hong Gu

Received: 13 March 2010 / Accepted: 30 July 2010  
© Springer-Verlag 2010

**Abstract** Starch paste viscosity plays an important role in estimating the cooking, eating, and processing quality of rice. The inheritance of starch paste viscosity in glutinous rice remains undefined. In the present study, 118 glutinous rice accessions were collected, and the genotypes of 17 starch synthesis-related genes (SSRG) were analyzed by using 43 gene-specific molecular markers. Association analysis indicated that 10 of 17 SSRGs were involved in controlling the rapid visco analyzer (RVA) profile parameters. Among these, the *PUL* gene was identified to play an important role in control of peak viscosity (PKV), hot paste viscosity (HPV), cool paste viscosity (CPV), breakdown viscosity (BDV), peak time (PeT), and paste temperature (PaT) in glutinous rice. Other SSRGs involved only a few RVA profile parameters. Furthermore, interactions between SSRGs were found being responsible for PeT, PaT, and BDV. Some of the RVA parameters, including PKV, HPV, CPV, CSV, and PaT, were mainly governed by single

SSRG, whereas other parameters, such as BDV, SBV, and PeT, were controlled by a few SSRGs, functioning cooperatively. Further, three near-isogenic lines (NIL) of a *japonica* glutinous cv. Suyunuo as genetic background, with *PUL*, *SSIII-1*, and *SSIII-2* alleles replaced with those of *indica* cv. Guichao 2, were employed to verify the genetic effects of the various genes, and the results were consistent with those obtained from the association analysis. These findings indicated that starch paste viscosity in glutinous rice had a complex genetic system, and the *PUL* gene played an important role in determining the RVA profile parameters in glutinous rice. These results provide important information for potentially improving the quality of glutinous rice.

**Keywords** Glutinous rice · Association analysis · Starch synthesis related genes · RVA profile parameters

## Introduction

Rice is one of the most important crops as it provides the staple food for half of the world's population, and high yield and good quality are two priorities in rice production. Due to the successful utilization of the semi-dwarf gene *sd-1* and heterosis technology, grain yield has been dramatically improved over the past several decades. However, far more improvements in the grain quality are required to meet the demand of consumers in rice producing areas.

Cooking and eating quality is widely considered as a major criteria for the grain quality, mainly determined by three physical and chemical indices, amylose content (AC) (Juliano 1985), gel consistency (GC) (Cagampang et al. 1973), and gelatinization temperature (GT) (Little et al.

---

Communicated by J. Snape.

---

C.-J. Yan and Z.-X. Tian contributed equally to this work.

---

C.-J. Yan · Y.-W. Fang · Y.-C. Yang · J. Li · S.-Y. Zeng ·  
S.-L. Gu · C.-W. Xu · S.-Z. Tang · M.-H. Gu (✉)  
Jiangsu Key Laboratory of Crop Genetics and Physiology,  
Key Laboratory of Plant Functional Genomics, Ministry  
of Education of China, Agricultural College of Yangzhou  
University, 12 East Wenhui Road, Jiangsu 225009,  
People's Republic of China  
e-mail: gumh@yzu.edu.cn

Z.-X. Tian  
Institute of Genetics and Developmental Biology,  
Chinese Academy of Science, Beijing 100101, China

1958). In the past decades, many researches focused on the genetic natures of these physicochemical indices to identify the genes controlling grain quality (He et al. 1999; Tan et al. 1999; Bao et al. 2000a, 2002; Lanceras et al. 2000; Septiningsih et al. 2003; Aluko et al. 2004; Tian et al. 2005; Fan et al. 2005; Wang et al. 2007). The accumulated results showed that AC and GC were largely determined by *Wx* gene, locating on chromosome 6 and encoding the granule-bound starch synthase (He et al. 1999; Tan et al. 1999; Septiningsih et al. 2003; Fan et al. 2005; Wang et al. 2007; Mikami et al. 2008). Similarly, GT was mainly controlled by *ALK* (or *SSII-3*) gene (He et al. 1999), encoding soluble starch synthase IIa (Umemoto et al. 2002; Umemoto and Aoki 2005; Gao et al. 2003).

Association analysis is a powerful tool for studying genetic loci involved in the inheritance of complex traits (Abdurakhmonov and Abdurkarimov 2008; Yu and Buckler 2006; Remington et al. 2001), and it has been successfully exploited in plant molecular genetics (Whitt et al. 2002; Wilson et al. 2004; Aranzana et al. 2005; Cockram et al. 2008). Through association analysis approach, a complex network controlling eating and cooking quality was identified, revealing that different characteristics were controlled by different gene combinations, and the genetic diversity of SSRGs greatly accounted for the varied grain quality (Tian et al. 2009).

Starch is composed of two forms, amylose and amylopectin. The fact that the cooking and eating quality varied among cultivars with similar AC suggests that the structure of amylopectin also has an effect in determining the physical and chemical properties (Juliano 1985). Moreover, the true genetic effects of other starch synthesis-related genes (SSRGs) are usually masked by the *Wx* gene (Bao et al. 2002), and the interactions between *Wx* and other SSRGs were often detected (He et al. 2006; Wu et al. 2006). The exact genetic effects of other SSRGs in shaping the rice grain quality are unclear yet.

Glutinous rice lacks starch amylose, which constitutes up to 30% of the total starch in non-glutinous rice endosperm. Glutinous rice is generally reserved for use in festival foods and desserts, although it also serves as the staple food in upland regions of Southeast Asia (Roder et al. 1996). Currently, it is widely used as an industry resource due to its specific property of lacking amylose in endosperm. Similar to common *indicaljaponica* cultivars, the glutinous rice varieties vary dramatically with respect to the eating and cooking quality, but the mechanism underlying the eating and cooking quality in glutinous rice is still unclear. Most researches on rice grain quality have been conducted on non-glutinous rice (Bao and Xia 1999; Bao et al. 2000a, b; Liu et al. 2004; Ge et al. 2005; He et al. 2006; Wang et al. 2007), only a few being on glutinous rice (Han et al. 2004; Bao et al.

2006). Therefore, we urgently need to reveal the genetic basis of eating and cooking quality in glutinous rice, and to provide the foundation for quality improvement in this food product.

It is well known that there are multiple isoforms of four classes of enzymes involved in starch synthesis, including ADP-glucose pyrophosphorylase (AGPase), starch synthase (SS), starch branching enzymes (SBE) and starch debranching enzymes (DBE), and each enzyme functions distinctly (Myers et al. 2000; Nakamura 2002). In the absence of the *Wx* gene, the glutinous rice has little or no amylose and is mainly composed of amylopectin, enabling us to study the genetic effects of SSRGs in glutinous rice. In previous studies, the effects of only a few genes were investigated in glutinous rice (Bao et al. 2006; Han et al. 2004). However, ~20 genes encoding four classes of enzymes are involved in amylopectin biosynthesis, and each gene plays a distinct role (Nakamura 2002). Therefore, until now, the genetic basis of glutinous rice quality remains obscure. Such questions as which gene(s) is the key factor(s) in determining the glutinous rice quality, and how they function remain to be addressed. The answers to these questions will be helpful in elucidating the mechanism of glutinous rice grain cooking and eating quality.

Recently, the RVA profile of starch paste viscosity was employed to evaluate rice eating and cooking quality (Juliano 1996). The RVA profile has become increasingly popular for investigating the viscosity property, since it requires only a small sample size and the procedure is easy to perform (Bao and Xia 1999). Here, we used a population of 118 glutinous rice varieties to investigate the genetic effects of 17 SSRGs involved in the starch biosynthesis system on formation of the grain RVA profile through association analysis. Additionally, the genes controlling grain RVA profile parameters without the effect of *Wx* were identified and verified with near-isogenic SSRG lines.

## Materials and methods

### Plant materials

In the present study, we collected 118 glutinous rice accessions from China and International Rice Research Institute (Table 1). Most of the accessions are *indica* and *japonica* types in *O. sativa*, while eight accessions belong to *O. glaberrima*. These glutinous rice accessions were planted in the experimental farm of Yangzhou University and Hainan in 2006 and 2007 in the rice growing season. At maturity, the seeds from each accession were harvested for RVA profile measurement.

**Table 1** Glutinous rice varieties employed in the present study

Code	Accession	Description	Origin	Code	Accession	Description	Origin
1	Hongmangxiangjingnuo	<i>Japonica</i>	China	60	Yangnuo No. 2	<i>Japonica</i>	China
2	Xiangjingnuo	<i>Japonica</i>	China	61	Jingnuo No. 6	<i>Indica</i>	China
3	Xiangjingnuo-1	<i>Japonica</i>	China	62	Lixiaonuo	<i>Indica</i>	China
4	Xiangmangnuo	<i>Japonica</i>	China	63	Baotanu	<i>Indica</i>	China
5	Xiangzhunuo	<i>Japonica</i>	China	64	Xiangjingnuo 103	<i>Japonica</i>	China
6	Xiangzhunuo-xuan	<i>Japonica</i>	China	65	Yangnuo No. 5	<i>Japonica</i>	China
7	Wujinxiangnuo	<i>Japonica</i>	China	66	Xiannuo 201	<i>Indica</i>	China
8	Jinggunuo	<i>Japonica</i>	China	67	Yangzhou No. 4	<i>Japonica</i>	China
9	Hongkenuo	<i>Japonica</i>	China	68	Gehuxiangnuo	<i>Japonica</i>	China
10	Hongkenuo-1	<i>Japonica</i>	China	69	Jiangzhouxiangnuo	<i>Japonica</i>	China
11	Baikenuo	<i>Japonica</i>	China	70	Yangxiannuo 32-2	<i>Indica</i>	China
12	Baikenuo-1	<i>Japonica</i>	China	71	Yandao No. 5	<i>Indica</i>	China
13	Baikenuo-2	<i>Japonica</i>	China	72	Xiangjingnuo 259	<i>Japonica</i>	China
14	Hongmangnuo	<i>Japonica</i>	China	73	Huinuo	<i>Japonica</i>	China
15	Hongmangnuo-1	<i>Japonica</i>	China	74	99-25	<i>Japonica</i>	China
16	Hongmangnuo-2	<i>Japonica</i>	China	75	Shuangfeng No. 4	<i>Japonica</i>	China
17	Jintannuo	<i>Japonica</i>	China	76	Yaxueno	<i>Indica</i>	China
18	Jintannuo-1	<i>Japonica</i>	China	77	Baixiangnuo	<i>Japonica</i>	China
19	Liyangnuo	<i>Japonica</i>	China	78	Shangnongxiangnuo	<i>Japonica</i>	China
20	Liyangnuo-1	<i>Japonica</i>	China	79	Taihunuo	<i>Japonica</i>	China
21	Wannuodao	<i>Japonica</i>	China	80	Shiwuyenuo	<i>Japonica</i>	China
22	Wannuodao-1	<i>Japonica</i>	China	81	Jingnuo 96103	<i>Japonica</i>	China
23	Shuangjiangqinnuodao	<i>Japonica</i>	China	82	Guanglingxiangnuo	<i>Japonica</i>	China
24	Xueliqinnuodao	<i>Japonica</i>	China	83	T002	<i>Japonica</i>	China
25	Xicainuo	<i>Japonica</i>	China	84	Changsiruanzhan	<i>Indica</i>	China
26	Changjingnuo	<i>Japonica</i>	China	85	Hongzaonuo	<i>Indica</i>	China
27	Shuijinnuo	<i>Japonica</i>	China	86	G3-66	<i>Japonica</i>	China
28	Huangjingnuo	<i>Japonica</i>	China	87	Cungunuo	<i>Indica</i>	China
29	Putaoonuo	<i>Japonica</i>	China	88	TAPOL	<i>Indica</i>	Philippines
30	Putaoonuo-1	<i>Japonica</i>	China	89	BLACK GORA S.N. 109	<i>Indica</i>	India
31	Zhongqiuputaoonuo	<i>Japonica</i>	China	90	CHALBYEO	<i>Japonica</i>	Korea
32	Huajiaonuo	<i>Japonica</i>	China	91	SHIKOKU MOCHI	<i>Japonica</i>	Japan
33	Huakenuo	<i>Japonica</i>	China	92	SHAN KIU JU	<i>Japonica</i>	China
34	Bainuodao	<i>Japonica</i>	China	93	KININPOL	<i>Japonica</i>	Philippines
35	Zhuzhongnuo	<i>Japonica</i>	China	94	MINAMI-HATA MOCHI	<i>Japonica</i>	Japan
36	Yangnuodao	<i>Japonica</i>	China	95	ARC 10283	<i>Javanica</i>	India
37	Yangnuodao-1	<i>Japonica</i>	China	96	CHIBA-A-HO-MOCHI	<i>Javanica</i>	Brazil
38	Zinuo	<i>Japonica</i>	China	97	HSINCHU GLUTTNOUS	<i>Javanica</i>	Brazil
39	Hongmangnuo-3	<i>Japonica</i>	China	98	PULUT PUTEH	<i>Javanica</i>	Malaysia
40	Hongmangnuo-4	<i>Japonica</i>	China	99	DAENG MAFAI	Intermediate	Thailand
41	Jintainuo	<i>Japonica</i>	China	100	ANANDI	Intermediate	Nepal
42	Huangnuo	<i>Japonica</i>	China	101	MIMIDAM	Intermediate	Bangladesh
43	Hongkenuo-2	<i>Japonica</i>	China	102	TININTA(MALAGKIT)	Intermediate	Philippines
44	Baikenuo-3	<i>Japonica</i>	China	103	AIBAIKOU	Intermediate	China
45	Jiangyinnuo	<i>Japonica</i>	China	104	KATUPA-AI	Intermediate	Taiwan
46	Zaonuodao	<i>Japonica</i>	China	105	KETAN TRENGGALEK	Intermediate	Indonesia
47	Chushunuo	<i>Japonica</i>	China	106	ARC 13163	Intermediate	India
48	Shixingnuodao	<i>Japonica</i>	China	107	HAWM OM	<i>O. glaberima</i>	Thailand

**Table 1** continued

Code	Accession	Description	Origin	Code	Accession	Description	Origin
49	Zhendao No2	<i>Japonica</i>	China	108	SUWEON 311	<i>O. glaberima</i>	Korea
50	Yannuo 98-9	<i>Japonica</i>	China	109	TAICHUNG SEN GLUTINOUS 1	<i>O. glaberima</i>	Taiwan
51	Xuenuo	<i>Indica</i>	China	110	TAICHUNG SEN GLUTINOUS 2	<i>O. glaberima</i>	Taiwan
52	Xinxiangnuo	<i>Japonica</i>	China	111	HIRAKAWA OKUTE	<i>O. glaberima</i>	Japan
53	57697	<i>Indica</i>	China	112	PULUTAN(PUTI)	<i>O. glaberima</i>	Philippines
54	535	<i>Indica</i>	China	113	XIANG N004	<i>O. glaberima</i>	China
55	Xiaomakenuo	<i>Indica</i>	China	114	HUANG SI NOO	<i>O. glaberima</i>	China
56	Bendixiangnuo	<i>Indica</i>	China	115	Makenuo	<i>Japonica</i>	China
57	Baiainuo	<i>Indica</i>	China	116	Zhenzhunuo	<i>Japonica</i>	China
58	3401	<i>Indica</i>	China	117	Heijienuo	<i>Japonica</i>	China
59	Henuo	<i>Indica</i>	China	118	Liutiaonuo	<i>Japonica</i>	China

No. 88–114 were introduced from Genetic Resources Center of IRRI

In order to verify the genetic effects of the identified genes obtained from association analysis approach, the near-isogenic lines (NIL) of the corresponding genes were developed. In the process of development of NIL, a typical *indica* cultivar Guichao 2 (inferior quality) and a glutinous *japonica* cultivar Suyunuo (good quality), were selected as donor and recipient, respectively, to produce a cross, and then continuously backcrossed for eight generations by molecular marker-aided selection. Finally, a series of NILs were obtained, and the investigation of their genetic background was conducted with 80 SSR markers covering the entire genome and 17 markers specific to starch synthesis genes. These NILs were also planted in the experimental farm of Yangzhou University in the rice growing season in 2008, and the seeds were harvested for RVA profile assessment.

#### RVA profile measurement

The RVA profiles were measured on a rapid visco analyser (RVA) (Model No. RVA-3D, Newport Scientific, Sydney, Australia), according to the Standard Method AACCC61-02 given by the American Association of Cereal Chemists (2000). Briefly, approximately 3 g rice flour was mixed with 25 ml water; a paddle was placed in the canister and rotated at 960 rpm for 10 s to disperse the rice sample. The viscosity was evaluated using a constant paddle rotation of 160 rpm. The sequential temperature curve for a 12.5 min test was as follows: (1) incubate at 50°C for 1.0 min; (2) increase to 95°C; (3) keep at 95°C for 1.4 min; (4) cool down to 50°C; and (5) hold at 50°C for 1.4 min. Viscosity values were recorded in centipose (cp). Starch viscosity characteristics included the following original components: peak viscosity (PKV), hot paste viscosity (HPV), and cool paste viscosity (CPV). Three secondary parameters including breakdown

(BDV), setback (SBV), and consistency (CSV) were calculated based on the original data:  $BDV = PKV - HPV$ ,  $SBV = CPV - PKV$  and  $CSV = CPV - HPV$ . In addition, pasting temperature (temperature of the initial viscosity increase, PaT) and pasting time (time of the initial viscosity increase, PeT) were also recorded.

#### Molecular marker development and assessment

In order to determine the diversity of SSRGs at the genomic level, a total of 13 representative cultivars, including 6 *japonica* cv. Nipponbare, Chunjiang 06, Wuyujing 7, Suyunuo, Taihunuo, Jiangzhouxiangnuo, and 7 *indica* cv. 93-11, 9308, Minghui 63, Longtefu, Guichao 2, TN 1 and Zhenshan 97, were selected and 17 SSRGs (including *AGPiso*, *AGP<sub>sma</sub>*, *AGP<sub>lar</sub>*, *GBSS II*, *SSI*, *SSII-1*, *SSII-2*, *SSII-3*, *SSIII-1*, *SSIII-2*, *SSIV-1*, *SSIV-2*, *SBE1*, *SBE3*, *SBE4*, *ISA*, and *PUL*) from each cultivars were then cloned and sequenced (Tian et al. 2009). Based on the genomic diversity of SSRGs, 43 STS/CAPs markers (Table 2) were developed and subsequently employed for genotyping 118 glutinous accessions (Tian et al. 2010).

#### Population structure and association analysis

The population structure was evaluated by using the STRUCTURE program (Pritchard et al. 2000) with 45 simple sequence repeats (SSR) distributed on 12 chromosomes in rice, including RM259, RM5, RM128, RM14, RM211, RM475, RM263, RM525, RM16, RM251, RM489, RM520, RM514, RM335, RM471, RM252, RM255, RM122, RM289, RM587, RM412, RM242, RM205, RM528, RM585, RM11, RM180, RM234, RM336, RM264, RM308, RM223, RM316, RM566, OSR28, RM333, RM216, RM258, RM474, RM286, RM332, RM441, RM17, RM101, and

**Table 2** Gene-tagged markers for SSRGs

Genes name	Markers name	Primer sequence	Markers type
<i>AGPlar</i>	AGPlar M1	[F] CGTTCAGGTTTCAGGCAATCA [R] GGAAGGGTGGTGTATGTGGAG	STS
	AGPlar M2	[F] GCGTGAAGTGAACATCCATCT [R] GGTTCAGCCTTCAGGTCAG	CAPS ( <i>Tsp45I</i> <sup>a</sup> )
<i>AGPiso</i>	AGPiso M2	[F] CAATCGCTGCCATCGGTTG [R] TTCCACATCGTTAGGTACACG	STS
	AGPiso M3	[F] TGGAATGGGAAGTCTATTATTGG [R] TCCCAACCTCTACCTTCAAATG	CAPS ( <i>EcoRI</i> )
<i>AGPsmA</i>	AGPsmA M1	[F] TCTATTCTCAGCCCTCCAACC [R] GTGTGTTTAGAGGTGCTTTTCG	STS
	AGPsmA M2	[F] TACGCTATGCTCTTGAAAC [R] TATCTTCCAGTAACCATCA	STS
<i>GBSSII</i>	GBSSII M1	[F] TTGCTGCGAATTATCTGCG [R] ACCTCCTCCCCTCTTTGC	STS
<i>SSI</i>	SSI M2	[F] CTTCTATCCATTCTTAATCCCA [R] ATGCTATTGATGTTAAGAGGGC	STS
	SSI M3	[F] GACCCACCTCGCTATCTGTTG [R] GGAAACACCAGACATCAACCAG	CAPS ( <i>ApaI</i> )
<i>SSII-1</i>	SSII-1 M1	[F] CACCCACCGTTCTACTATGC [R] TCCATAGTTTCATTGAGATTGCTC	STS
	SSII-1 M2	[F] CAAGTTGGTGACGATAGTGATGA [R] AACAGAGCCTCCATTACCTTTAC	CAPS ( <i>AgeI</i> )
	SSII-1 M3	[F] AGAGATCAAATCGTGGAAC [R] TGGAGTGAAGTAGTGAAT	STS
	SSII-1 M4	[F] ATCTTTAGACGATTAGCG [R] AAGTCACAAGTAGAAGGG	STS
<i>SSII-2</i>	SSII-2 M1	[F] AGATTTGAACTCAGGACTTGGTG [R] TCTATGGGCTCTATCCTTACTAGG	STS
	SSII-2 M3	[F]ACAGTATGTTTGCCTCAGCG [R] GTAAATCCACCCAGCCAGTC	STS
<i>SSII-3</i>	SSII-3 M1	[F] CCAATACCGTAAACTAGCGACTATG [R] TACAGGTAGAATGGCAGTGGTG	STS
	SSII-3 M2	[F] GGTTCTCGGTGAAGATGGC [R] GTGGTCCCAGCTGAGGTCC	CAPS ( <i>BanII</i> )
	SSII-3 M3	[F] AACTGACTCATAACGGATAACG [R] CACGCACGAACGGAAACC	CAPS ( <i>NheI</i> )
<i>SSIII-1</i>	SSIII-1 M1	[F] AAGAAGGGAAGGGAGTCAGC [R] GCCATCTCCATTGCCAGC	SSR
	SSIII-1 M2	[F] CAAGCAATGATTCAGGCACA [R] GGAGACAGGAGCAAAAGGC	CAPS ( <i>EcoRI</i> )
	SSIII-1 M3	[F] CAAATCAACTGTAAGTGCTGGAG [R] GAGAACGGAGAAAATGGCAT	CAPS ( <i>NdeI</i> )
<i>SSIII-2</i>	SSIII-2 M1	[F] AAGTCCTTCGGCTTACTATTCC [R] GGAGAAGGAACATAACAGGGAC	CAPS ( <i>XbaI</i> )
	SSIII-2 M2	[F] GAACCTGTGCCTTAAGCTGACTG [R] GGAATAGTAAGCCGAAGGACTT	STS
<i>SSIV-1</i>	SSIV-1 M1	[F] CATTGTGTCTTGAAGTCTGTGCT [R] CGATGGGTTAGTGCTGTGG	CAPS ( <i>NdeI</i> )

Table 2 continued

Genes name	Markers name	Primer sequence	Markers type
SSIV-2	SSIV-2 M1	[F] CTTCTGATTGATGGTTGGTTGC [R] GGAAGAATAATCTCTACTAGGTGGC	CAPS ( <i>Sph</i> I)
	SSIV-2 M2	[F] TTCCCTTGGTGGTGCGTG [R] TAAAGCGTTCGACAGTA	STS
	SSIV-2 M3	[F] TCAAGTATGGTTTACCTATG [R] TTTCCCAATGACTTCTAA	CAPS ( <i>Eco</i> 72I)
SBE1	SBE1 M1	[F] TGCTACATAACACGCATACAAAAGT [R] AGACAAAAGCGAAAAGGTAATGAG	STS
	SBE1 M2	[F] GTGGGGAAAACAAGTAAGTCTG [R] AGTTCCATCAGAAGAATCAGGG	STS
	SBE1 M3	[F] GGAAATGGGAGTCGCC [R] CGAAGAAACCACGCTCA	STS
	SBE1 M4	[F] ATTGTTGCTGAAGATGTTT [R] ACGGTTGATGGTAGGTG	CAPS ( <i>Taq</i> I)
SBE3	SBE3 M2	[F] GTGGGGTTCTCAACTAGC [R] CATCAGCATTGTTAGGCAG	STS
SBE4	SBE4 M1	[F] CACCAATTATATTAGCGTGCTCC [R] CGTGGCTCTTGGCTCTCTTG	STS
	SBE4 M2	[F] CCATCACCTCAAATACATCACTC [R] AACTGGAATGCCCTTAGG	STS
ISA	ISA M1	[F] ATAGATGCTAATGTGATGTGGC [R] TGGTATAGGCACAACCGTAGA	STS
	ISA M2	[F] ACAAGCACACGACACCTA [R] CAACAAAACCAAATCATT	CAPS ( <i>Hind</i> III)
	ISA M3	[F] TGTGGGAATACCTTCAACTG [R] ATAAAACCCCTTACAGGCTTG	STS
PUL	PUL M1	[F] AGAGAAGGAGAAAAGAAGTGGAGAC [R] GTCCAAACTGAATCACTCAATCG	STS
	PUL M2	[F] CCACCATTAAGCATCATCAAC [R] AGTTGTTATATTTTAGGATGGATGG	STS
	PUL M3	[F] CTGTATGGACTGAGTAGTCGATGG [R] TGAGCCTCATCTGCCAGAGT	STS
	PUL M4	[F]TACACCATCTCACTACCA [R] GCAACATCTAAAACACCAA	STS
	PUL M5	[F]ATTGGCATTGTGAAGTTTC [R] CAATCTTGGTTTTATCCTG	STS
	PUL M6	[F]ATTTAACTGTATGGACTGAG [R] GATACCAACCAAACAAGA	STS

<sup>a</sup> Represents the enzymes used for CAPS

RM519. The number of subgroups was determined to be two based on an admixture and linkage model, which agreed with prior population information. The resulting Q-values were obtained from the STRUCTURE program. The relative kinship (K) matrix was calculated using the software package SPAGeDi (Hardy and Vekemans 2002). Association analysis followed the unified

mixed model previously reported (Yu et al. 2005), using SAS 9.0.

Statistical analysis

A *t* test program in SPSS 10.0 was employed to determine the presence of significant differences between the RVA

profile parameters of Suyunuo and those of NIL-*PUL*, NIL-*SSIII-1*, and NIL-*SSIII-2*.


## Results

### Phenotypic variations within the glutinous rice accessions

We planted 118 glutinous rice accessions in Yangzhou and Hainan in 2006 and 2007, respectively, and seeds from each accession were harvested for assessments of amylose content and RVA parameters. Due to its expected significant effect on RVA parameters, the amylose contents for 118 glutinous accessions were first assessed. The results showed that the amylose contents of all samples were less than 3% (data not shown), indicating that the 118 accessions in the present study were truly glutinous rice. The RVA measurements were then performed. It was found that

the RVA data collected over the 2 years were very similar, and significant correlations were observed (data not shown). Therefore, we averaged the 2-year RVA data for the following analysis. The accessions were ordered according to PKV value, and eight RVA parameters of the 118 accessions are shown in Fig. 1 and Table 3.

As shown in Fig. 1, most of the RVA parameters, including PKV, HPV, CPV, BDV, and CSV, of *japonica* accessions were generally less than those of the *indica* accessions. For example, there were 77 accessions whose PKV values were <1,700 cp; of which, 69 accessions belonged to the *japonica* type, whereas, only seven accessions belonged to the intermediate type, and one was *O. glaberrima*. In the 41 accessions whose PKV were bigger than 1,700 cp, there were 20 *indica*, 10 *japonica*, 3 *java-nica* and 3 intermediate types. As for SBV, the reverse trend was observed. This result suggested that the genes in control of the RVA profile parameters have differentiated, at least between the *indica/japonica* subspecies. Moreover,



**Fig. 1** RVA profile parameters of 118 glutinous rice accessions. **a–h** PKV, HPV, BDV, CPV, SBV, CSV, PeT, and PaT, respectively. The *x*-axis denotes the accession code, which is ordered based on the PKV value

**Table 3** Statistical analysis of RVA profile parameters in 118 glutinous rice accessions

Parameters	Mean	Standard deviation	Minimum	Maximum
PKV (cp)	1,570	517.4	610	3,084
HPV (cp)	731.7	401.34	139	1,875
BDV (cp)	838.6	281.1	285	1,712
CPV (cp)	958.5	531.4	242	3,363
SBV (cp)	-611.8	347.9	-1,516	952
CSV (cp)	230.4	186.6	98	1,488
PeT (min)	3.8	0.6	3	6.6
PaT (°C)	71.9	2.9	64.1	82.35

The results were obtained by averaging 2 years' data in 2006 and 2007

most of the RVA parameters, such as PKV, HPV, CPV, SBV, BDV, and CSV, varied widely among the 118 glutinous accessions (Table 3). For example, the PKV value ranged from 868 to 2,711 cp, with an average value of 1,570 cp. By contrast, PaT and PeT had relatively small variations, with PaT in most of accessions being very close to about 70°C, few exceptions at 80°C. As for PeT, most of the accessions were ~3.6 min, while only 18 accessions (15%) were ~5 min (Table 3). These results suggest that, in the present study, the 118 glutinous accessions are representative in terms of rice grain quality and are qualified for genetic analysis of RVA parameters.

It also should be noted that several accessions have dramatically higher or lower values in CPV, SBV, CSV, and PeT, distinct from the other accessions, including

No. 26 (Putanuo, *japonica*), No. 31 (Wujinxiangnuo, *japonica*), No. 59 (Ketan Trenggalek, intermediate), No. 63 (Bainuodao, *japonica*), No. 76 (Hongmangnuo, *japonica*), No. 86 (Suweon 311, *O. glaberrima*), and No. 105 (Tininta, intermediate) (Fig. 1d–g). The existence of distinct glutinous rice accessions implies that in glutinous rice germplasm, the genes underlying the eating and cooking quality have differentiated dramatically, although these genes are unknown yet. These distinct accessions may be helpful in the development of special varieties for industry.

#### Correlation among RVA parameters

To explore the relationship among eight RVA profile parameters, the pairwise correlation analysis was conducted, and the correlation coefficients among eight RVA profile parameters were summarized in Table 4. Interestingly, the significant correlations were found between almost any two parameters; and only three pairwise correlations between BDV and HPV, CPV, CSV did not reach the significant level (Table 4). The result suggested that the eight RVA profile parameters in the 118 glutinous accessions were interdependent, similar to the results by Wang et al. (2007).

Although significant correlations among the eight RVA parameters were observed, many exceptions also existed. For example, in general, PKV was positively correlated with HPV; however, the accession Liutianuo had high PKV (2,446 cp), but its HPV was relatively low (734 cp). In addition, CPV in most of the glutinous accessions were <1,500 cp, while some accessions, such as Bainuodao, Liyangnuo, Suweon 311 and Yangxiannuo 32-2, exhibited

**Table 4** Correlation analysis among eight RVA profile parameters

	PKV	HPV	BDV	CPV	SBV	CSV	PeT
HPV	0.84217 <0.0001						
BDV	0.63821 <0.0001	0.12237 <b>0.0605</b>					
CPV	0.78015 <0.0001	0.96536 <0.0001	0.05766 <b>0.3779</b>				
SBV	-0.2953 <0.0001	0.22216 0.0006	-0.86081 <0.0001	0.36727 <0.0001			
CSV	0.41498 <0.0001	0.60785 <0.0001	-0.10404 <b>0.1109</b>	0.77712 <0.0001	0.56982 <0.0001		
PeT	0.41478 <0.0001	0.74557 <0.0001	-0.30104 <0.0001	0.8559 <0.0001	0.69043 <0.0001	0.83403 <0.0001	
PaT	0.43673 <0.0001	0.68504 <0.0001	-0.17421 0.0073	0.65111 <0.0001	0.34504 <0.0001	0.38751 <0.0001	0.66857 <0.0001

The number in the upper line is the correlation coefficient, and the number in the lower line indicates the corresponding probability  $P > 0.05$  showed in bold



bigger CPV, more than 1,500 cp. These results primarily suggested that different genetic mechanisms are responsible for the different RVA profile parameters.

#### Association analysis

In association analysis, the population structure usually plays an important role in identifying target gene loci. Therefore, in order to eliminate the influence of population structure effects on association analysis, we analyzed the population structure by using the STRUCTURE software based on the genotypes of 45 SSR markers covering the entire rice genome. The results showed that 118 glutinous accessions could be divided into two groups (Fig. 2), which fit well with their species-specific properties (*indica* vs. *japonica*). The  $Q$  value of each accession generated from population structure analysis was used in the following association analysis.

The association analysis result was summarized in Table 5. As shown, three original parameters, PKV, HPV, and CPV were interestingly found to be mainly controlled by the *PUL* gene, suggesting that starch (amylopectin) viscosity was mainly affected by this gene in glutinous rice. By contrast, two secondary parameters, BDV and SBV, several genes were identified to be responsible for their variations. For BDV, four genes, including *AGPlar*, *PUL*, *SSI*, and *SSIV-1*, were shown to be responsible for its variation in 118 glutinous accessions. Seven genes, including *AGPlar1*, *SBE1*, *SBE3*, *ISA*, *SSII-1*, *SSII-2*, and *SSIV-1*, were found to contribute to the SBV variation in the glutinous rice population, of which, *SSIV-1* had the largest effect according to the  $F$  value. However, for another secondary parameter, CSV, only one gene, *SBE1*, was found to be responsible for its variation.


As for PeT parameter, which reflects the time of the initial viscosity increase, three genes, *SSII-3*, *PUL*, and *AGPlar*, were identified. Among these, *SSII-3* had the largest genetic effect with an additive effect of  $0.82^{\circ}\text{C}$ , apparently functioning as a key factor in determining the PeT parameter. However, only the *PUL* gene was found to associate with PaT variation, which represents the temperature of the initial increase in starch viscosity.


To comprehensively understand the genetic mechanisms underlying the RVA profile parameters, the interactions among 17 genes were analyzed (Fig. 3). The result showed that complex genetic interactions existed for controlling the BDV parameter; the network contained not only four identified genes (*PUL*, *SSI*, *AGPlar*, and *SSIV-1*) when a single gene was considered in association analysis, but also three additional genes, including *ISA*, *SBE1*, and *SSII-2*, which were not detected to be independently responsible for BDV variations. The fact that the genetic network involved four classes of enzymes, AGPase, SBE, DBE, and SSS, indicated that there was a very complex genetic system underlying BDV performance. Similarly, interactions between *SSII-3* and *PUL* as well as between *SSII-3* and *SSIII-2* were also identified to contribute PaT parameter. Furthermore, an interaction between *PUL* and *SSIII-2* was also found to affect the PeT parameter. Apart from

**Table 5** Results of association analysis between SSRG and RVA profile parameters

Trait	SSRG	Estimate	$F$ value	$P$ value
PKV	<i>PUL</i>	442.02	38.07	1.03E-06
HPV	<i>PUL</i>	295.09	38.69	3.14E-06
BDV	<i>AGPlar</i>	264.25	5.58	1.10E-03
	<i>PUL</i>	138.06	8.33	6.92E-03
	<i>SSI</i>	223.78	14.59	2.54E-04
	<i>SSIV-1</i>	238.97	7.41	6.69E-03
CPV	<i>PUL</i>	266.68	26.31	3.93E-04
SBV	<i>AGPlar</i>	385.48	9.00	1.53E-04
	<i>SBE1</i>	281.45	8.56	4.47E-03
	<i>SBE3</i>	258.70	7.51	7.45E-03
	<i>ISA</i>	288.56	7.49	7.07E-03
	<i>SSII-1</i>	213.50	6.60	1.07E-02
	<i>SSII-2</i>	397.35	11.12	1.08E-03
	<i>SSIV-1</i>	405.97	13.64	3.17E-04
CSV	<i>SBE1</i>	104.41	6.98	8.72E-04
PeT	<i>AGPlar</i>	0.51	4.89	7.14E-03
	<i>PUL</i>	0.29	8.75	4.45E-03
	<i>SSII-3</i>	0.82	4.81	1.69E-03
PaT	<i>PUL</i>	1.83	12.34	2.34E-03

**Fig. 2** Population structure analysis. The results of structure analysis indicated that two groups ( $K = 2$ ) exist in this population, and the  $Q$  value of each individual are shown





**Fig. 3** A diagram showing the genetic interactions controlling BDV, PaT, and PeT parameters. The *arrows* between any two genes indicate the existence of an interaction between them

BDV, PaT, and PeT, we failed to detect any interactions between the other five RVA profile parameters.

Taken together, our analysis revealed that in 17 SSRGs, the *PUL* gene was shown to be heavily involved in the regulation of most of the RVA profile parameters in glutinous rice; the complex genetic systems associated with RVA profile parameters were controlled through both individual SSRGs as well as through interactions between SSRGs.

#### Verification of genetic effects of SSRG through NILs

To validate the results of the association analysis, three near-isogenic lines, NIL-*PUL*, NIL-*SSIII-1*, and NIL-*SSIII-2*, were employed to evaluate the variations of RVA profile parameters. These three NILs were generated by marker-assisted selection toward target genes *PUL*, *SSIII-1*, and *SSIII-2* in the progeny of backcrossed population derived from a cross of Suyunuo (a *japonica* glutinous landrace, recipient) × Guichao 2 (an *indica* cultivar, donor). The result was shown in Table 6. When compared to the recurrent parent Suyunuo, significant changes occurred in PKV, HPV, BDV, CPV, and SBV of NIL-*PUL*, whereas CSV, PeT and PaT did not alter significantly. This result was basically consistent with the results obtained in the association analysis, confirming that the *PUL* gene plays an important role in shaping the eating and cooking quality in glutinous rice. However, no any significant changes in RVA parameters were observed in NIL-*SSIII-1* when compared to its recurrent parent Suyunuo. In fact, in the association analysis, *SSIII-1* showed no effects on eight RVA parameters, suggesting that there was no functional differentiation at the *SSIII-1* locus, at least between Guichao 2 and Suyunuo, or that it may be compensated by other SSS. As for NIL-*SSIII-2*, the significant changes in PeT and PaT were detected compared to the recurrent parent Suyunuo, and other RVA profile parameters were

**Table 6** Comparison of RVA profile parameters between recurrent parent and NILs

Materials	Sample size	PKV (cp)	HPV (cp)	BDV (cp)	CPV (cp)	CSV (cp)	SBV (cp)	PeT (min)	PaT (°C)
Suyunuo	4	1,037.25 ± 119.76	346.50 ± 52.74	690.75 ± 73.95	491.5 ± 66.38	145 ± 14.63	-545.75 ± 59.97	3.50 ± 0.03	71.05 ± 0.44
NIL- <i>SSIII-1</i>	10	1,030.70 ± 169.74	358.70 ± 59.26	672.00 ± 112.04	513.00 ± 68.38	154.3 ± 12.09	-517.70 ± 109.21	3.47 ± 0.09	70.71 ± 0.81
NIL- <i>PUL</i>	4	1,278 ± 53.63*	470.50 ± 30.69**	807.50 ± 29.15*	625.25 ± 39.48*	154.75 ± 0.49	-652.75 ± 25.32*	3.53 ± 0.05	70.7 ± 0.34
NIL- <i>SSIII-2</i>	4	1,184.75 ± 116.13	364.75 ± 27.50	820.00 ± 97.36	513.50 ± 33.97	148.75 ± 10.72	-617.25 ± 87.18	3.36 ± 0.04**	69.35 ± 0.46**

\* Significant at 5% level, \*\*significant at 1% level

not altered significantly; meanwhile, the *SSIII-2* was found to be incorporated into the SSRG interactions for regulating PaT and PeT in association analysis. Therefore, the finding based on three NILs was basically consistent with those from association analysis with respect to genetic effects of SSRGs.

## Discussion

Glutinous rice is generally reserved for use in festival foods and desserts, and it also serves as the staple food in the upland regions of Southeast Asia (Roder et al. 1996). Now, the glutinous rice also is widely used as an industry resource. However, little efforts have been made to reveal the genetic mechanism of the starch paste viscosity in the glutinous rice. In the present study, the genetic mechanism of starch viscosity in glutinous rice was analyzed by using the association analysis method, and the results provided an overview of the genetic basis of the starch viscosity profile in glutinous rice. Such finding will potentially benefit future efforts to improve the quality of glutinous rice, and perhaps even of the common *indica* and *japonica* rice.

The paste viscosity profile is considered as a very important predictor of the eating, cooking, and processing quality characteristics of common *indica/japonica* rice (Juliano 1985; Shu et al. 1998). Thus, information on the genes affecting paste viscosity parameters will facilitate our efforts to improve rice grain quality, although the relationship between paste viscosity profile and eating and cooking quality in glutinous rice remains unclear yet. In previous studies, Bao et al. (2006) developed several microsatellite markers for the *SSI* and *SBE1* loci on the basis of sequence diversity, and the relationship between the genotypes on *SBE1* and *SSI* loci and starch thermal and retrogradation properties, pasting properties, and swelling volume in 56 glutinous rice accessions were analyzed. The results showed that accessions with different genotypes on the two loci exhibited different starch properties, suggesting that alleles on *SBE1* and *SSI* loci have functionally differentiated. Meanwhile, Han et al. (2004) analyzed the effects of the *SBE1* and *SBE3* genes on the paste viscosity in 40 glutinous rice lines, and the result indicated that polymorphisms on both *SBE1* and *SBE3* loci accounted for ~70% of the observed variations in both hot and cool viscosities, and for ~40% of the observed variations in peak viscosity and consistency. However, in the two aforementioned studies, the population sizes were relatively small, and a few of SSRGs were considered. In particular, the effect of population structure on the gene effects analysis was neglected, and these shortcomings may have led to inaccurate conclusions about the gene functions. In the present study, we employed a larger

population (118 glutinous accessions), and most of the SSRGs (17) and population structures were considered as a whole to analyze the genetic behavior of starch paste viscosity parameters; for these reasons, our conclusions may be more reasonable.

In the past decades, more researches on rice grain quality were performed in non-glutinous rice, in which the role of amylopectin in eating and cooking quality is, at least in part, masked by amylose (Bao et al. 2002), whereas in glutinous rice which contains only amylopectin, the eating and cooking quality would be predominantly affected by the properties of this polysaccharide. Hence, some genes with minor effects may possibly be overlooked in non-glutinous rice. Gravois and Webb (1997) previously analyzed the genetic behavior of rice viscosity, and their results showed that PKV, HPV, and CPV were controlled by one major gene (*Wx*). Bao et al. (2000a) mapped the QTL for RVA parameters using a doubled haploid population derived from a cross between an *indica* variety Zai-Ye-Qing 8 and a *japonica* variety Jing-Xi 17, and a total of 20 QTL for six parameters of the RVA profiles were identified at least in one location; meanwhile, only the *Wx* gene was detected significantly in both environments for five traits (HPV, CPV, BDV, CSV, and SBV), indicating that the RVA profiles were mainly governed by the *Wx* locus, although some minor QTL were also identified to be responsible for the RVA profiles (Bao et al. 2000b). These results were further confirmed by Wang et al. (2007), in which a recombinant inbred population derived from a cross between Zhenshan 97 and Delong 208 was used to analyze the genetic basis of RVA profiles. Thus, it is clear that the *Wx* gene is a key determinant in the control of RVA profile parameters in common *indica* and *japonica*. However, in glutinous rice, loss-of-function mutation occurring in the *Wx* gene resulted in the failure of amylose synthesis (Wanchana et al. 2003). Although the *Wx* gene itself does not function in glutinous rice, RVA profile parameters vary widely in glutinous rice accessions. Therefore, it seems that the genetic information gained from analysis of rice grain quality in non-glutinous rice cannot be directly used for glutinous rice analysis and improvement. In the present experiment, the association analysis results showed that most of the SSRGs (10/17) were responsible for the paste viscosity profiles, and the *PUL* gene was shown to play an important role in the control of most of the RVA profile parameters, except for CSV and SBV, in glutinous rice. Unfortunately, the genetic effect of the *PUL* gene was not detected in studies by both Wang et al. (2007) and Bao et al. (2000b). This result may be caused by two factors. First, the genetic populations in both of these studies were derived from a common *indica/japonica* cross, in which *Wx* gene works normally and thus, amylose is produced in the rice endosperm. Current

knowledge indicates that the *Wx* gene functions as a major factor in determining RVA profile parameters, and the effect of *PUL* gene can be possibly masked by *Wx*. In fact, when the *PUL* allele in Guichao 2 was substituted with that of Suyunuo, no RVA profile parameters altered significantly when compared to those of Guichao 2 (Wu et al. 2006). Second, the discrepancy between our study and the two reports mentioned above could be that in the Zhai-Ye-Qing 8/Jing-Xi 17 and Zhenshan 97/Delong 208 derived populations, there was no functional differentiation on the *PUL* locus between the two parents, thus leading to the failure of identification of its genetic effect. Taken together, the data suggest that in order to improve grain quality of common *indica* and *japonica* cultivars, we should focus on the selection of favorable alleles on the *Wx* locus in order to regulate the amylose content; however, some amylopectin synthesis related genes, such as *SSII-3*, *PUL*, and *SSIII-2*, which have influence on determining the fine structure of amylopectin, should not be neglected.

It is generally accepted that ADPase, SSS, SBE, and DBE were involved in starch biosynthesis in plants (Nakamura 2002). There were two types of DBE, isoamylase (ISA) and pullulanase (PUL, also known as limit dextrinase or R-enzyme), classified on the basis of their sequence similarity and substrate specificity. Kubo et al. (1999) reported that both ISA and PUL were involved in amylopectin biosynthesis in rice endosperm; they also presumed that ISA played a predominant role in amylopectin synthesis, but PUL was also essential or compensated for the role of ISA in the formation of the amylopectin multiple-cluster structure. The analysis of three PUL-deficient mutants indicated that the short chain ( $DP \leq 13$ ) of amylopectin in *PUL* mutants was increased, and the average chain length of B2-3 chains was  $\sim 3$  residues longer compared with that of the wild-type (Fujita et al. 2009). Therefore, it is obvious that *PUL* plays a role in determining the fine structure of amylopectin. In our present study, *PUL* gene was found to be a principal determinant for the variation in RVA profile parameters in glutinous rice, consistent with the results obtained in the analysis of *PUL* mutants.

Both association analysis and NIL analysis indicated that *SSIII-2* was one of regulators of the PaT and PeT parameters, suggesting that this gene indeed plays a role in formation of the fine structure of amylopectin. A previous study showed that the loss-of-function of *SSIII-2* resulted in the reduction of the amylopectin B2–B4 chains with degree of polymerization ( $DP \geq 30$ ) by  $\sim 60\%$  of the wild-type values, strongly suggesting *SSIII-2* functions in the elongation of amylopectin B2 to B4 chains (Fujita et al. 2007). Hence, it can be speculated that the chains with  $DP \geq 30$  in amylopectin contributes to the eating and cooking quality, which can be reflected in the PeT and PaT parameters.

Based on the results of our study, it therefore can be deduced that functional differentiations occurred in most of the SSRGs, especially at the *PUL*, *SSII-3*, and *SSIII-2* loci, and different alleles at the *PUL* and *SSIII-2* loci have different genetic effects on RVA profile parameters. In fact, the exploitation of multiple alleles have been widely conducted at the *Wx* and *SSII-3* loci, which were shown as major factors in the control of AC and GT, respectively (Umamoto et al. 2002; He et al. 2006; Wang et al. 2007). To date, at least five alleles at the *Wx* locus were identified in Asian rice germplasm (Mikami et al. 2008), which provides the foundation to explain the continuous variation of AC in Asian rice landraces. Similarly, there are two alleles at the *SSII-3* locus identified in Asian cultivars, and the diversity of the *SSII-3* locus results in the different amylopectin structure and starch quality between *japonica* and *indica* varieties (Umamoto et al. 2002; Nakamura et al. 2005; Bao et al. 2006; Tian et al. 2009), being reflected in the difference of the GT among these varieties. Moreover, in the present study, Guichao 2 and Suyunuo belong to typical *indica* and *japonica* types, respectively, and great sequence diversities were identified on each SSRGs (Tian et al. 2009). The functional differentiations that occurred in a few SSRGs were deduced on the basis of the performances of their NILs. Therefore, with more detailed studies performed, increasing information on allelic diversification of SSRGs loci would be obtained, enabling us to purposefully select a favorable allele on each locus by molecular marker-assisted selection and consequently to develop new varieties to meet the demand of rice consumers.

Although most of the RVA profile parameters were found to be controlled by a few SSRGs, two important parameters, BDV and SBV, previously shown to be closely related to eating and cooking quality (Bao and Xia 1999), were found here to be governed by at least four SSRGs, and various interactions were also found to be in the control of BDV, PaT, and PeT. This result implies that, manipulation of a few SSRGs cannot achieve the goal of improvement of rice grain quality. Moreover, the interactions between SSRGs remain unclear, and more NILs and transgenic lines for SSRGs are needed for evaluating the gene functions and understanding the complex genetic network responsible for starch quality.

**Acknowledgments** We are grateful to Mr. Xie Yulin (Taihu Institute of Agricultural Science, Jiangsu, China) for providing some materials. This study was financially supported by the Ministry of Science and Technology (Grant No. 2005CB120804 and 2006AA10Z118), the National Natural Science Foundation (Grant Nos. 30530470, 30771323 and 30871501), Fok Ying Tung Education Foundation (101030) and the Jiangsu Province Government (Grant No. 08KJA210002) of China.

## References

- Abdurakhmonov IY, Abdurakimov A (2008) Application of association mapping to understanding the genetic diversity of plant germplasm resources. *Int J Plant Genomics*. doi:10.1155/2008/574927
- Aluko G, Martinez C, Tohme J, Castano C, Bergman C, Oard JH (2004) QTL mapping of grain quality traits from the interspecific cross *Oryza sativa* × *O. glaberrima*. *Theor Appl Genet* 109: 630–639
- American Association of Cereal Chemists (2000) Approved methods for the AACCC, 10th edn. Method 61–01 (amylograph method for milled rice) and Method 61–02 (determination of the pasting properties of rice with rapid visco analyzer). The Association, St. Paul
- Aranzana MJ, Kim S, Zhao K, Bakker E, Horton M, Jakob K, Lister C, Molitor J, Shindo C, Tang C, Toomajian C, Traw B, Zheng H, Bergelson J, Marjoram P, Dean, Nordborg M (2005) Genome-wide association mapping in *Arabidopsis* identifies previously known flowering time and pathogen resistance genes. *PLoS Genet* 1(5):e60
- Bao JS, Xia YW (1999) Genetic control of paste viscosity characteristics in *indica* rice (*Oryza sativa* L.). *Theor Appl Genet* 98:1120–1124
- Bao JS, He P, Li SG, Xia YW, Chen Y, Zhu LH (2000a) Comparative mapping quantitative trait loci controlling the cooking and eating quality of rice. *Sci Agric Sin* 33:8–13 (in Chinese with English abstract)
- Bao JS, Zheng XW, Xia YW, He P, Shu QY, Lu X, Chen Y, Zhu LH (2000b) QTL mapping for the paste viscosity characteristics in rice (*Oryza sativa* L.). *Theor Appl Genet* 100:280–284
- Bao JS, Wu YR, Hu B, Wu P, Cui HR, Shu QY (2002) QTL for rice grain quality based on a DH population derived from parents with similar apparent amylose content. *Euphytica* 128:317–324
- Bao JS, Corke H, Sun M (2006) Microsatellites, single nucleotide polymorphisms and a sequence tagged site in starch-synthesizing genes in relation to starch physicochemical properties in nonwaxy rice (*Oryza sativa* L.). *Theor Appl Genet* 113:1185–1196
- Cagampang GB, Perez CM, Juliano BO (1973) A gel consistency test for eating quality in rice. *J Sci Food Agric* 24:1589–1594
- Cockram J, White J, Leigh FJ, Lea FJ, Chiapparino E, Laurie DA, Mackay IJ, Powell W, O'Sullivan DM (2008) Association mapping of partitioning loci in barley. *BMC Genet* 9:16. doi:10.1186/1471-2156-9-16
- Fan CC, Yu XQ, Xing YZ, Xu CG, Luo LJ, Zhang QF (2005) The main effects, epistatic effects and environmental interactions of QTLs on the cooking and eating quality of rice in a doubled-haploid line population. *Theor Appl Genet* 110:1445–1452
- Fujita N, Yoshida M, Kondo T, Saito K, Utsumi Y, Tokunaga T, Nishi A, Satoh H, Park JH, Jane JL, Miyao A, Hirochika H, Nakamura Y (2007) Characterization of SSIIIa-deficient mutants of rice: the function of SSIIIa and pleiotropic effects by SSIIIa deficiency in the rice endosperm. *Plant Physiol* 144:2009–2023
- Fujita N, Toyosawa Y, Utsumi Y, Higuchi T, Hanashiro I, Ikegami A, Akuzawa S, Yoshida M, Mori A, Inomata K, Itoh R, Miyao A, Hirochika H, Satoh H, Nakamura Y (2009) Characterization of pullulanase (PUL)-deficient mutants of rice (*Oryza sativa* L.) and the function of PUL on starch biosynthesis in the developing rice endosperm. *J Exp Bot* 60(3):1009–1023
- Gao ZY, Zeng DL, Cui X, Zhou YH, Yan M, Huang D, Li JY, Qian Q (2003) Map-based cloning of the *ALK* gene, which controls the GT of rice. *Sci China (Ser C)* 46:661–668
- Ge XJ, Xing YZ, Xu CG, He YQ (2005) QTL analysis of cooked rice grain elongation, volume expansion, and water absorption using a recombinant inbred population. *Plant Breed* 124:121–126
- Gravois KA, Webb BD (1997) Inheritance of long grain rice amylograph viscosity characteristics. *Euphytica* 97:25–29
- Han YP, Xu ML, Liu XY, Yan CJ, Korban SS, Chen XL, Gu MH (2004) Genes coding for starch branching enzymes are major contributors to starch viscosity characteristics in waxy rice (*Oryza sativa* L.). *Plant Sci* 166:357–364
- Hardy OJ, Vekemans X (2002) SPAGeDi: a versatile computer program to analyze spatial genetic structure at the individual or population levels. *Mol Ecol Notes* 2:618–620
- He P, Li SG, Qian Q, Ma YQ, Li JZ, Wang WM, Chen Y, Zhu LH (1999) Genetic analysis of grain quality. *Theor Appl Genet* 98:502–508
- He Y, Han YP, Jiang L, Xu CW, Lu JF, Xu ML (2006) Functional analysis of starch-synthesis genes in determining rice eating and cooking qualities. *Mol Breed* 18:277–290
- Juliano BO (1985) Criteria and test for rice grain quality. In: Juliano BO (ed) *Rice chemistry and technology*. American Association of Cereal Chemists Inc., St. Paul, pp 443–513
- Juliano BO (1996) Rice quality screening with the Rapid Visco Analyser. In: Walker CE, Hazelton JL (eds) *Applications of the rapid visco analyser*. Newport Scientific, Sydney, pp 19–24
- Kubo A, Fujita N, Harada K, Matsuda T, Satoh H, Nakamura Y (1999) The starch debranching enzymes isoamylase and pullulanase are both involved in amylopectin biosynthesis in rice endosperm. *Plant Physiol* 121:399–409
- Lanceras JC, Huang ZL, Naivikul O, Vanavichit A, Ruanjaichon V, Tragoonrun S (2000) Mapping of genes for cooking and eating qualities in Thai jasmine rice (KDML105). *DNA Res* 7:93–101
- Little RR, Hilder GB, Dawson EH (1958) Differential effect of dilute alkali on 25 varieties of milled white rice. *Cereal Chem* 35:111–126
- Liu XY, Gu MH, Han YP, Ji Q, Lu JF, Gu SL, Zhang R, Li X, Chen JM, Korban SS, Xu ML (2004) Developing gene-tagged molecular markers for functional analysis of starch-synthesizing genes in rice (*Oryza sativa* L.). *Euphytica* 135:345–353
- Mikami I, Uwatoko N, Ikeda Y, Yamaguchi J, Hirano HY, Suzuki Y, Sano Y (2008) Allelic diversification at the wx locus in landraces of Asian rice. *Theor Appl Genet* 116:979–989
- Myers AM, Morell MK, James MG, Ball SG (2000) Recent progress toward understanding biosynthesis of the amylopectin crystal. *Plant Physiol* 122:989–997
- Nakamura Y (2002) Towards a better understanding of the metabolic system for amylopectin biosynthesis in plants: rice endosperm as a model tissue. *Plant Cell Physiol* 43(7):718–725
- Nakamura Y, Francisco PB Jr, Hosaka Y, Sato A, Sawada T, Kubo A, Fujita N (2005) Essential amino acids of starch synthase IIa differentiate amylopectin structure and starch quality between *japonica* and *indica* rice varieties. *Plant Mol Biol* 58:213–227
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* 155(2):945–959
- Remington DL, Thornsberry JM, Matsuoka Y, Wilson LM, Whitt SR, Doebley J, Kresovich S, Goodman MM, Buckler ES (2001) Structure of linkage disequilibrium and phenotypic associations in the maize genome. *Proc Natl Acad Sci USA* 98:11479–11484
- Roder W, Keoboulapha B, Vannalath K, Phouaravanh B (1996) Glutinous rice and its importance for hill farmers in Laos. *Econ Bot* 50:401–408
- Septiningsih EM, Trijatmiko KR, Moeljopawiro S, McCouch SR (2003) Identification of quantitative trait loci for grain quality in an advanced backcross population derived from the *Oryza sativa* variety IR64 and the wild relative *O. rufipogon*. *Theor Appl Genet* 107:1433–1441
- Shu QY, Wu DX, Xia YW, Gao MW, McClung A (1998) Relationship between RVA profile character and eating quality in *Oryza sativa* L. *Sci Agric Sin* 31:25–29

- Tan YF, Li JX, Yu SB, Xing YZ, Xu CG, Zhang QF (1999) The three important traits for cooking and eating quality of rice grains are controlled by a single locus in an elite rice hybrid, Shanyou 63. *Theor Appl Genet* 99:642–648
- Tian R, Jiang GH, Shen LH, Wang LQ, He YQ (2005) Mapping quantitative trait loci underlying the cooking and eating quality of rice using a DH population. *Mol Breed* 15:117–124
- Tian ZX, Qian Q, Liu QQ, Yan MX, Liu XF, Yan CJ, Liu GF, Gao ZY, Tang SZ, Zeng DL, Wang YH, Yu JM, Gu MH, Li JY (2009) Allelic diversity in rice starch biosynthesis pathway leads to a diverse array of rice eating and cooking qualities. *Proc Natl Acad Sci USA* 106:21760–21765
- Tian ZX, Yan CJ, Qian Q, Yan S, Xie HL, Wang F, Xu JF, Liu GF, Wang YH, Liu QQ, Tang SZ, Li JY, Gu MH (2010) Development of gene-tagged molecular markers for starch synthesis related genes in rice. *Chinese Sci Bull* (in press)
- Umemoto T, Aoki N (2005) Single-nucleotide polymorphisms in rice starch synthase IIa that alter starch gelatinization and starch association of the enzyme. *Funct Plant Biol* 32:763–768
- Umemoto T, Yano M, Satoh H, Shomura A, Nakamura Y (2002) Mapping of a gene responsible for the difference in amylopectin structure between *japonica*-type and *indica*-type rice varieties. *Theor Appl Genet* 104:1–8
- Wanchana S, Toojinda T, Tragoonrun S, Vanavichit A (2003) Duplicated coding sequence in the *waxy* allele of tropical glutinous rice (*Oryza sativa* L.). *Plant Sci* 165:1193–1199
- Wang LQ, Liu WJ, Xu Y, He YQ, Luo LJ, Xing YZ, Xu CG, Zhang QF (2007) Genetic basis of 17 traits and viscosity parameters characterizing the eating and cooking quality of rice grain. *Theor Appl Genet* 115:463–476
- Whitt SR, Wilson LM, Tenaillon MI, Gaut BS, Buckler ES (2002) Genetic diversity and selection in the maize starch pathway. *Proc Natl Acad Sci USA* 99:12959–12962
- Wilson LM, Whitt R, Ibañez AM, Rocheford TR, Goodman MM, Buckler ES (2004) Dissection of maize kernel composition and starch production by candidate gene association. *Plant Cell* 16:2719–2733
- Wu HK, Liang GH, Gu YJ, Shan LL, Wang F, Han YP, Gu MH (2006) The effect of the starch-synthesizing genes on RVA profile characteristics in rice (*Oryza sativa* L.). *Acta Agro Sin* 32(11):1597–1603 (in Chinese with English abstract)
- Yu J, Buckler ES (2006) Genetic association mapping and genome organization of maize. *Curr Opin Biotechnol* 17:155–160
- Yu JM, Pressoir G, Briggs WH, Bi IV, Yamasaki M, Doebley JF, McMullen MD, Gaut BS, Nielsen DM, Holland JB, Kresovich S, Buckler ES (2005) An unified mixed-model method for association mapping that accounts for multiple levels of relatedness. *Nat Genet* 38:203–208